

THE SPIN STRUCTURE FUNCTION g_2 and ASYMMETRY A_2 FROM SLAC E155X


STEPHEN ROCK
UNIVERSITY OF MASSACHUSETTS

REPRESENTING THE E155X COLLABORATION

Sept, 2002

- EXPERIMENT
- SOME THEORY
- g_2 AND A_2 FOR PROTON/DEUTERON
- Q^2 DEPENDENCE OF g_2 AND A_2
- TWIST-3 MATRIX ELEMENT d_2
- SUM RULES

Polarized Deep Inelastic Scattering




$$A_{\perp} = \frac{\sigma^{\downarrow\leftarrow} - \sigma^{\uparrow\leftarrow}}{\sigma^{\downarrow\leftarrow} + \sigma^{\uparrow\leftarrow}} = f_k E' \sin(\theta) \left[\mathbf{g}_1(x, Q^2) + \frac{2E}{\nu} \mathbf{g}_2(x, Q^2) \right]$$

$$A_{\parallel} = \frac{\sigma^{\uparrow\downarrow} - \sigma^{\uparrow\uparrow}}{\sigma^{\uparrow\downarrow} + \sigma^{\uparrow\uparrow}} = f_k \left\{ \mathbf{g}_1(x, Q^2) [E + E' \cos(\theta)] - \frac{Q^2}{\nu} \mathbf{g}_2(x, Q^2) \right\}$$

\mathbf{g}_1 and \mathbf{g}_2 are the polarized structure functions.

- \mathbf{A}_{\parallel} is primarily sensitive to \mathbf{g}_1
- \mathbf{A}_{\perp} is more sensitive to \mathbf{g}_2
- f_k includes contribution from kinematics and unpolarized structure functions


E155 Spectrometers



EXPERIMENTAL OVERVIEW

- **SLAC ELECTRON BEAM:**
 - POLARIZATION $P_b = 0.83$
 - ENERGY $E = 29.1$ or $E = 32.3$
- **TRANSVERSELY POLARIZED TARGETS:**
 - 1 K, 5 T, D.N.P
 - NH_3 (PROTONS) $\langle P_t \rangle = 0.70$
 - LiD (DEUTERONS) $\langle P_t \rangle = 0.22$
 - RADIATION DAMAGE
- **SCATTERED ELECTRONS**
 - 2.75° , 5.5° , AND 10.5°
 - CERENKOV AND LEAD GLASS FOR PARTICLE I.D.
 - HODOSCOPES FOR TRACKS
($dP/P \approx 0.03$)

KINEMATICS



THE A_{\perp} ASYMMETRY

$$\tilde{A}_{\perp} = \frac{1}{f_{RC}} \left[\frac{C_1}{fP_t} \left(\left(\frac{N_L - N_R}{N_L + N_R} \right) \frac{1}{P_b} + A_{EW} \right) + C_2 \right] + A_{RC}$$

- N_L AND N_R ARE RATES FOR L AND R BEAM HELICITY, CORRECTED FOR PAIR-SYMMETRIC CONTRIBUTIONS (FEW PERCENT) AND π MIS-IDENTIFIED AS ELECTRONS (FEW PERCENT)
- $C_1 \approx 1$ CORRECTS FOR POLARIZED ^{15}N or Li
- ADDITIONAL CORRECTION FOR ^7Li and LiH CONTAMINATION PLANE OF ELECTRON SCATTERING.
- f IS DILUTION FACTOR (≈ 0.16 PROTON)
- A_{EW} : ELECTROWEAK ASYMMETRY ($\approx 8 \times 10^{-5} Q^2$)
- f_{RC} AND A_{RC} : RADIATIVE CORRECTIONS
 - f_{RC} Mostly from Elastic and Quasi-elastic tail: Increases Statistical Error
 - A_{RC} Mostly From Resonance and DIS

DETERMINATION OF g_2 and A_2

- WE USE MEASURED \tilde{A}_\perp and E155 g_1 FIT TO OBTAIN g_2 USING:

$$g_2 = \frac{yF_1}{2E'(\cos \Theta - \cos \alpha)} \left[\tilde{A}_\perp \nu \frac{(1 + \epsilon R)}{1 - \epsilon} - \frac{g_1}{F_1} [E \cos \alpha + E' \cos \Theta] \right]$$

- WE CAN ALSO EXTRACT VIRTUAL PHOTON-NUCLEON ASYMMETRY

$$A_2(x, Q^2) = \frac{2\sigma_{TL}}{\sigma_T^{1/2} + \sigma_T^{3/2}} = \gamma \frac{g_1(x, Q^2) + g_2(x, Q^2)}{F_1(x, Q^2)}$$

- COMPARE TO POSITIVITY LIMIT

$$A_2 < \sqrt{R(x, Q^2)}$$

- COMPARE TO SOFFER LIMIT

$$A_2 < \sqrt{R(1 + A_1)/2}$$

SYSTEMATIC ERRORS

A_{\perp} ERRORS

- RADIATIVE CORRECTIONS
 - Used Many Different Models of A_{\perp}
 - Resonance, twist-3, Quasi-elastic ..
- MULTIPLICATIVE ERR: 5.1% (p); 6.2% (d)
 - TARGET POL: 1.7% (p); 4.5% (d)
 - DILUTION FACTOR: 3.0% (p); 2.1% (d)
 - BEAM POLARIZATION: 3.7%

g_2 ERRORS

- g_1 : 5% INCLUDES MODEL ERRORS
- POLARIZATION ANGLE
- F_2 : 2%
- R

SIMPLE FIT USED TO CALCULATE g_2^{WW}

$$Q^2 > 1$$

$$g_1^p/F_1^p = x^{.700}(.817+1.014x-1.489x^2)(1-.04/Q^2)$$

$$g_1^n/F_1^n = x^{-.335}(-.013-.330x+.761x^2)(1+.13/Q^2)$$

Q^2 DEPENDENCE

Proton -0.04 ± 0.06

Neutron 0.13 ± 0.45

OTHER FITS USED: INCLUDED
IN SYSTEMATIC ERROR ON

g_1

g₂(x, Q²) STRUCTURE FUNCTION

- In general

twist-2
↓

twist-2
↓

twist-3
↓


$$\begin{aligned}
 g_2(x, Q^2) &= g_2^{\text{WW}}(x, Q^2) - \int_x^1 \frac{\partial}{\partial y} \left(\frac{m}{M} h_T(y, Q^2) + \xi(y, Q^2) \right) \frac{dy}{y} \\
 &= g_2^{\text{WW}}(x, Q^2) + \bar{g}_2(x, Q^2)
 \end{aligned}$$

- Wandzura-Wilczek g_2 expression (twist-2 only)


$$g_2^{\text{WW}}(x, Q^2) = -g_1(x, Q^2) + \int_x^1 \frac{dy}{y} g_1(y, Q^2)$$

- \bar{g}_2 includes quark-gluon correlations (ξ) inside the nucleon
- Twist-2 term (quark transverse spin distribution h_T) in \bar{g}_2 suppressed by $\frac{m}{M} \ll 1$

Q^2 -DEPENDENCE OF xg_2




RESULTS FOR Q^2 -AVERAGED xg_2



- - - - WW - . - - Stratmann Song
 - - - - Weigel - - - - Wakamatsu
 ◇ E143 □ E155

χ^2/df agreement with $g_2^{WW} = 3.1(p) \quad 1.2(d)$

RESULTS FOR Q^2 -AVERAGED A_2



◇ E143 E155 SMC
- - - $\sqrt{R(1 + A_1)/2}$

Operator Product Expansion

(OPE) For g_1 and g_2

OPE USED IN QCD TO SEPARATE THE PHYSICS INTO A PERTURBATIVE PIECE (easily treated) AND A NON-PERTURBATIVE PIECE (unknown matrix elements).

$$\Gamma_1^{(n)} = \int_0^1 x^n g_1(x, Q^2) dx = \frac{a_n}{2}, \quad n = 0, 2, 4, \dots$$

$$\Gamma_2^{(n)} = \int_0^1 x^n g_2(x, Q^2) dx = \frac{1}{2} \frac{n}{n+1} (d_n - a_n)$$

$$n=2,4,\dots$$

a_n are the twist-2 and d_n are the twist-3 matrix elements.

$$d_2 = \int_0^1 x^2 \left[2g_1(x, Q^2) + 3g_2(x, Q^2) \right] dx$$

$$= 3 \int_0^1 x^2 \overline{g_2}(x, Q^2) dx$$

Q² DEPENDENCE

- All the Integrals are at FIXED Q₀² while the Measurements have Q² and x correlated.

- For **g₁** integrals, we have 'evolved' to fixed Q² using Q² dependent fits to g₁/F₁

- For **g₂** we use:

$$g_2(x, Q_0^2) = \overline{g_2}(x, Q_m^2) + g_2^{WW}(x, Q_0^2)$$

and ASSUME that the Q² dependence of $\overline{g_2}(x, Q^2)$ is negligible.


- **d₂** depends logarithmically on Q².
- To calculate **g₂^{WW}** we utilize a global Q² dependent fit to **g₁/F₁**.
 - **g₁/F₁** has errors.
 - Statistical Errors on **g₂** Dominate
- Low x Extrapolation: Suppressed By x²
- High x Extrapolation: g₂ Very Small

THE TWIST-3 d_2 MATRIX ELEMENT

$$d_2 = 3 \int_0^1 x^2 [g_2(x, Q^2) - g_2^{WW}(x, Q^2)] dx$$

**E155X: $0.0025 \pm .0016 \pm .0010$ (proton)
 $0.0054 \pm .0023 \pm .0005$ (deuteron)**


AVERAGE: $0.0032 \pm .0017$ (p) $0.0079 \pm .0048$ (n)



QUARK MODEL WITH TRANSVERSE MOMENTUM

$$A_2F_1 \propto g_1 + g_2 = \int_x^{k_T^2 \max} dk_T^2 \tilde{h}\left(x + \frac{k_T^2}{xM^2}\right)$$

Roberts and Ross PROTON



BURKHARDT-COTTINGHAM SUM RULE

$$\int_0^1 g_2(x) dx = 0$$

- Derived from Virtual Compton Scattering Dispersion Relations.
- Does not follow from the OPE since $n = 0$.
- Validity depends on the lack of singularities for g_2 at $x = 0$.
 - Dramatic rise of g_2 at low x could invalidate the sum rule

$$\int_0^9 g_2(x) dx \text{ at } Q^2 = 5$$

E155X

AVERAGE

$$\text{P: } -0.044 \pm 0.008 \pm .003 \quad -0.042 \pm .008 \text{ (5 } \sigma)$$

$$\text{D: } -0.008 \pm 0.012 \pm .002 \quad -0.006 \pm .011$$

LOW x EXTRAPOLATION

$$\text{IF } g_2 = g_2^{WW} \text{ AT LOW } x$$

$$\int_0^x g_2^{WW}(y) dy = x \left[g_2^{WW}(x) + g_1(x) \right]$$

$$\int_0^9 = -0.022 \pm 0.008(\text{p}) \quad -0.002 \pm .011(\text{d})$$

EFREMOV-TERYAEV-LEADER SUM RULE

$$\int_0^1 \mathbf{x} [\mathbf{g}_1^V(\mathbf{x}) + 2\mathbf{g}_2^V(\mathbf{x})] d\mathbf{x} = 0$$

$$\text{IF } \Delta \mathbf{u}_{\text{sea}} = \Delta \mathbf{d}_{\text{sea}} \Rightarrow$$

$$\int_0^1 \mathbf{x} [\mathbf{g}_1^P(\mathbf{x}) + 2\mathbf{g}_2^P(\mathbf{x}) - \mathbf{g}_1^N(\mathbf{x}) - 2\mathbf{g}_2^N(\mathbf{x})] d\mathbf{x} = 0$$

E155X



AVERAGE

-0.013 ± .008 ± .002



-0.011 ± .008

**LOW x EXTRAPOLATION SUPPRESSED
BY x FACTOR**

TRANSVERSE PION ASYMMETRIES from P



TRANSVERSE PION ASYMMETRIES from D



g₂ SUMMARY

- PROTON g₂ IS POSITIVE FOR x < 0.1, NEGATIVE FOR x > 0.2.
- ROUGH AGREEMENT SEEN WITH SHAPE OF g₂^{WW} and BAG MODEL OF STRATMANN
- DISAGREES IN DETAIL WITH g₂^{WW}:
 $\chi^2/\text{dof} = 3.1$
- $A_2 < \sqrt{R(1 + A_1)}/2$
- PROTON A₂ IS POSITIVE FOR ALL x, A₂ → 0 FOR x → 0.
- d₂ SMALL
- ELT SUM RULE OK
- BURKHARDT-COTTINGHAM SUM RULE NOT OK FOR PROTON
- JLAB EXPERIMENTS ON g₂

E155/E155x Collaboration

P. Anthony, R. Arnold, T. Averett, H. Band, N. Benmouna, C. Berisso, H. Borel
P. Bosted, S. Bültmann, M. Buenerd, T. Chupp, S. Churchwell, G.R. Court
D. Crabb, D. Day, P. Decowski, P. DePietro, R. Erbacher, R. Erickson
A. Feltham, H. Fonvieille, E. Frlez, R. Gearhart, V. Ghazikhanian, J. Gomez
K. Griffioen, C. Harris, M.A. Houlden, E. W. Hughes, C. Hyde-Wright, G. Igo
S. Incerti, J. Jensen, J. Johnson, P. King, Yu. G. Kolomensky, S. Kuhn
R. Lindgren, R. Lombard-Nelsen, J. Marroncle, W. Meyer, J. McCarthy
P. McKee, D. McNulty, G. Mitchell, J. Mitchell, M. Olson, S. Penttila
G. Peterson, G. G. Petratos, R. Pitthan, D. Pocanic, R. Prepost, C. Prescott
L. M. Quin, B. Raue, D. Reyna, L.S. Rochester, S. Rock
O. Rondon-Aramayo, F. Sabatie, S. St. Lorant, I. Sick, L. Sorrell, T. Smith,
F. Staley L. M. Stuart, Z. Szalata, Y. Terrien, L. Todor, A. Tobias, T. Toole,
S. Trentalange, D. Walz, R. Welsh, F.R. Wesselmann, T. Wright, C. Young
B. Youngman, M. Zeier, H. Zhu, B. Zihlmann

American University

Universität Basel

University Blaise Pascal

Ruhr-Universität Bochum

California Inst. of Technology

University of California–Los Angeles

PNC Saclay

Florida International University

Kent State University

University of Liverpool

LANL

University of Massachusetts

University of Michigan

Old Dominion University

Smith College

SLAC

Stanford University

TJNAF

University of Virginia

College of William and Mary

University of Wisconsin